# Opioid Abuse in South Carolina: Current Perspectives








### **National Overdose Deaths**

Number of Deaths from Prescription Opioid Pain Relievers




Source: National Center for Health Statistics, CDC Wonder





### **National Overdose Deaths**


Number of Deaths from Heroin





Source: National Center for Health Statistics, CDC Wonder


1999


Drug-poisoning deaths are defined as having ICD-10 underlying cause-of-death codes X40-X44\* (unintentional)












































# Most Commonly Misused or Abused Types of Legal Drugs

# R4

### Opioid Pain Relievers

 drugs that contain active ingredient codeine, hydrocodone, and oxycodone.


Buprenorphine
Butorphanol (Stadol®)
Codeine
Fentanyl (Duragesic®
patch)
Hydrocodone (Vicodin®)
Hydromorphone
(Dilaudid®)
Meperidine (Demerol®)
Methadone

Morphine
Nalbuphine (Nubain®)
Oxycodone
(Percocet®/Percodan®)
Oxymorphone
Pentazocine (Talwin®)
Paregoric
Propoxyphene (Darvon®)



3 out of 4 people who used heroin in the

who used heroin in the past year misused opioids first



7 out of 10 people

who used heroin in the past year also misused opioids in the past year





Source: Jones, C.M., Heroin use and heroin use risk behaviors among nonmedical users of prescription opioid pain relievers - United States, 2002-2004 and 2008-2010. Drug Alcohol Depend. (2013).

# Addiction: Opioid Use Disorder





## Addiction: Opioid Use Disorder



- A disease of the brain
- Preventable
- Chronic

- Treatable
- Not Curable
- Sometimes Fatal

### Substance Use Disorders are Chronic Diseases

#### **DIABETES**

- Genetic predisposition
- Lifestyle choices are a factor in development of the disease
- Severity is variable
- There are diagnostic criteria
- Once diagnosed, you've got it

#### **ADDCITION**


- Genetic predisposition
- Lifestyle choices are a factor in development of the disease
- Severity is variable
- There are diagnostic criteria
- Once diagnosed, you've got it

# Medication-Assisted Treatment and Recovery for Opioid Use Disorders

A combination of behavioral therapy and medications (most commonly methadone, buprenorphine, and naltrexone)




### A note about environmental risk or protective factors...



## Recent Analysis and Action

- Inspector General's Report, May 2013
   "South Carolina Lacks a Statewide Prescription Drug Abuse Strategy"
- Executive Order No. 2014-22, March 2014
   Establishing Prescription Drug Abuse Prevention Council
- "State Plan to Prevent and Treat Prescription Drug Abuse"
   December 2014





Award from SAMHSA to Prevent Prescription Drug/Opioid Overdose-Related Deaths

# "The SC Overdose Prevention Project"


• **Funding amount:** \$3,192,772

Project Period: Five years










### **UNWANTED MEDICAL DISPOSAL SITES**

You can drop off unwanted prescription drugs at these locations year-round. Check with each location for hours of operation.





# State Targeted Response to the Opioid Crisis Grant (SAMHSA)

**Funding amount:** \$6,575,623

- Statewide Multi-Level Multimedia Campaign
- Healthcare Professional Training
  - Online Learning Collaborative
  - Academic Detailing
- SCRIPTS Integration With EHRs
- Offender Re-Entry Programming
- Financial Assistance for Treatment
- Recovery Programming
- Peer Support

### Save the Date

# Governor's Opioid Summit Sponsored by DAODAS

September 6<sup>th</sup> and September 7<sup>th</sup>

Columbia Metropolitan Convention Center

Sara Goldsby MSW, MPH sgoldsby@daodas.sc.gov 803-896-8371

